

# Upstream-migrating knickpoints: The dominant control on submarine channel evolution?

**Maarten S. Heijnen (1), Michael A. Clare (1), Matthieu J.B. Cartigny (2), Peter J. Talling (2), Sophie Hage (3), Gwyn Lintern (4), Cooper Stacey (4), Daniel R. Parsons (5), Justin K. Dix (3), John E. Hughes Clarke (6)**

(1) Marine Geosciences, National Oceanography Centre, Southampton, U.K., (2) Departments of Geography and Earth Sciences, University of Durham, U. K., (3) Ocean and Earth Sciences, National Oceanography Centre, University of Southampton, U.K., (4) Natural Resources Canada, Geological Survey of Canada, BC, Canada, (5) School of Environmental Sciences, University of Hull, U.K., (6) Earth Sciences, Center for Coastal & Ocean Mapping, University of New Hampshire, U.S.A.

[maarten.heijnen@noc.ac.uk](mailto:maarten.heijnen@noc.ac.uk)

Symposium Theme: Subaqueous mass movements in lakes, fjords and coastal areas

Keywords: Knickpoints, Fjord, Channel evolution, Time-lapse bathymetry, British Columbia

Submarine channel systems are the primary conduits for sediment transport to the deep sea. The submarine mass movements that pass through them are the volumetrically most significant sediment transport mechanisms on Earth. Submarine channel deposits preserve valuable archives of Earth history, while the associated mass movements pose a hazard to critical seafloor infrastructure. Therefore, understanding how submarine channels form and develop is important, particularly as this dictates where and how well deposits are preserved. It has been hypothesised that the development of meander bend cut offs due to outer bank erosion is a key control on submarine channel evolution. This hypothesis is based on river studies. Several recent studies show that flow in submarine channels is very different to that in fluvial settings, however. Other inferences have been made from the deposits left behind by past flows, scaled-down laboratory studies, and uncalibrated numerical models, but the key controls on the morphologic evolution of submarine channels remain elusive.

Here, we can reconcile these issues using unusually detailed time-lapse bathymetry surveys that record the morphologic evolution of a highly active submarine channel. Multiple seafloor surveys were performed in Bute Inlet, British Columbia over a ten-year period. We track the evolution of a 60 km-long submarine channel that is surveyed from its source to lobe at an unprecedented high spatial and temporal resolution. We first summarise the morphologic changes observed from sub-annual to decadal repeated surveys. Second, we classify the nature and quantify the scale of erosional and depositional patterns observed. Finally, we relate the observed morphologic changes to the processes that control channel evolution and discuss the implications of our findings for other submarine channel systems worldwide.

We find that upstream-migration of knickpoints (sharp and steep steps in the channel topography), is by far the most important process that drives channel evolution (Fig. 1). These features are an order of magnitude larger than the dune-scale crescentic bedforms that pervade much of the channel axis. Individual knickpoints can migrate several hundreds of metres upstream per year and can cause up to 30 metres depth of erosion over this entire migration distance. The influence of outer bend erosion is found to be negligible in comparison to knickpoint migration, with the latter capable of completely modifying both plan-view and cross-sectional morphology of the channel. High resolution bathymetry data reveals that similar knickpoints are also common in many other submarine channel systems worldwide, yet their potentially critical role in channel evolution has largely been ignored until now.

The submarine landslides observed offshore the Espírito Santo Basin reveal a variable internal complexity, relating to their remobilisation dynamics and the influence of salt structures on their origin. These aspects are relevant for the understanding of landslides and hazard assessment in similar salt-rich basins.




Figure 1: Slope map images of a section of the submarine channel in Bute Inlet in March 2008 (A) and October 2016 (B). The front of an area characterized by knickpoints has migrated upstream over several kilometres in the time between the surveys, and reshaped the channel. (C) Cross-sectional view through the channel at the location indicated in (A) and (B) shows the erosion associated with the upstream-migrating knickpoints.