

Accepted Manuscript

## *Geological Society, London, Special Publications*

### Revisiting the tsunamigenic volcanic flank-collapse of Fogo Island in the Cape Verdes, offshore West Africa

Rachel Barrett, Elodie Lebas, Ricardo Ramalho, Ingo Klaucke, Steffen Kutterolf, Andreas Klügel, Katja Lindhorst, Felix Gross & Sebastian Krastel

DOI: <https://doi.org/10.1144/SP500-2019-187>

Received 30 September 2019

Revised 13 November 2019

Accepted 25 November 2019

© 2019 The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by/4.0/>). Published by The Geological Society of London. Publishing disclaimer: [www.geolsoc.org.uk/pub\\_ethics](http://www.geolsoc.org.uk/pub_ethics)

To cite this article, please follow the guidance at <https://www.geolsoc.org.uk/onlinefirst#how-to-cite>

#### **Manuscript version: Accepted Manuscript**

This is a PDF of an unedited manuscript that has been accepted for publication. The manuscript will undergo copyediting, typesetting and correction before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the book series pertain.

Although reasonable efforts have been made to obtain all necessary permissions from third parties to include their copyrighted content within this article, their full citation and copyright line may not be present in this Accepted Manuscript version. Before using any content from this article, please refer to the Version of Record once published for full citation and copyright details, as permissions may be required.

## Revisiting the tsunamigenic volcanic flank-collapse of Fogo Island in the Cape Verdes, offshore West Africa

Rachel Barrett<sup>1\*</sup>, Elodie Lebas<sup>1</sup>, Ricardo Ramalho<sup>2,3,4,5</sup>, Ingo Klaucke<sup>6</sup>, Steffen Kutterolf<sup>6</sup>, Andreas Klügel<sup>7</sup>, Katja Lindhorst<sup>1</sup>, Felix Gross<sup>1,8</sup> & Sebastian Krastel<sup>1</sup>

<sup>1</sup>*Institute of Geosciences, Christian-Albrechts-University, Kiel, Germany*

<sup>2</sup>*Instituto Dom Luiz, Faculdade de Ciências da Universidade de Lisboa, Portugal*

<sup>3</sup>*Departamento de Geologia, Faculdade de Ciências da Universidade de Lisboa, Portugal*

<sup>4</sup>*School of Earth Sciences, University of Bristol, Bristol, UK*

<sup>5</sup>*Lamont-Doherty Earth Observatory of Columbia University, New York, USA*

<sup>6</sup>*GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel, Germany*

<sup>7</sup>*Department of Geosciences, University of Bremen, Bremen, Germany*

<sup>8</sup>*Kiel Marine Science, Christian-Albrechts-University, Kiel, Germany*

\*Correspondence ([rachel.barrett@ifg.uni-kiel.de](mailto:rachel.barrett@ifg.uni-kiel.de))

**Abstract:** Volcanic archipelagos are a source of numerous on- and offshore geohazards, including explosive eruptions and potentially tsunamigenic large-scale flank-collapses. Fogo Island in the southern Cape Verdes is one of the most active volcanoes in the world, making it both prone to collapse (as evidenced by the ca. 73 ka Monte Amarelo volcanic flank-collapse), and a source of widely-distributed tephra and volcanic material. The offshore distribution of the Monte Amarelo debris avalanche deposits and the surrounding volcanoclastic apron were previously mapped using only medium-resolution bathymetric data. Here, using recently acquired, higher resolution acoustic data, we revisit Fogo's flank-collapse, and find evidence suggesting that the deposition of hummocky volcanic debris originating from the failed eastern flank most likely triggered the contemporaneous, multi-phase failure of pre-existing seafloor sediments. Additionally, we identify, for the first time, multiple mass-transport deposits in the southern part of the volcanoclastic apron of Fogo and Santiago based on the presence of acoustically chaotic deposits in parametric echo sounder data and volcanoclastic turbiditic sands in recovered cores. These preliminary findings indicate a long and complex history of instability on the southern slopes of Fogo and suggest that Fogo may have experienced multiple flank collapses.

## Introduction

Volcanic archipelagos are home to numerous on- and offshore hazards such as changes in sea level, storms, volcanic eruptions, slope instabilities, large flank-collapses, and tsunamis (Casalbore 2018). These events can have devastating consequences for people, nature, and infrastructure, both onshore and offshore. A variety of internal and external factors can precondition and trigger flank instabilities. These include dyke and sill intrusions; volcanic eruptions and tremor; earthquakes; flank over-steepening; the weight of new volcanic material on the island flanks; weakening of the volcanic edifice by weathering and hydrothermal activity; and, for smaller landslides, by the effects of wave, wind and storm activity (e.g. Siebert 1984; Begét & Kienle 1992; Murray & Voight 1996; McGuire 1996, 2003; Tibaldi 2001; Cervelli *et al.* 2002; Casalbore *et al.* 2011, 2015; Gross *et al.* 2014; Clare *et al.* 2018). Such factors occur over timescales ranging from seconds to thousands of years, and are globally widespread (McGuire 1996; Blahůt *et al.* 2019).

The volume of volcanic flank-collapses varies significantly, but can be as large as thousands of cubic kilometres, e.g. in the Hawaiian archipelago (Moore *et al.* 1989). Many such events have volumes in the order of tens to hundreds of cubic kilometres, e.g. at Nisyros Volcano in the Aegean Sea (Tibaldi *et al.* 2008; Livanos *et al.* 2013); in the Lesser Antilles Arc (Lebas *et al.* 2011; Le Friant *et al.* 2015; Brunet *et al.* 2016); in the Canary Islands (Krastel *et al.* 2001; Masson *et al.* 2002; León *et al.* 2017); and in the Cape Verde Islands (Masson *et al.* 2008). However, even comparatively small-volume volcanic flank-collapses, such as the 0.22–0.3 km<sup>3</sup> Anak Krakatau flank-collapse in December 2018 (Grilli *et al.* 2019), may result in catastrophic tsunamis. The hazard potential of such flank-collapses is widely recognized, but the magnitude, and therefore hazard potential, of the tsunamis that can be triggered by flank-collapses is heavily debated (e.g. Moore & Moore 1984; Goff *et al.* 2014; McMurtry *et al.* 2004; Watt *et al.* 2012a; Ramalho *et al.* 2015; Paris *et al.* 2018).

Other factors that can contribute to the instability of slopes offshore are the deposition of centi- to decimetre-thick discrete layers of volcanic ash across a wide region and the presence of buried turbidites in the volcanic apron. Studies in lacustrine settings (e.g. Wiemer *et al.* 2015; Moernaut *et al.* 2019) and on active offshore margins (e.g. Harders *et al.* 2010; Lafuerza *et al.* 2014; Hornbach *et al.* 2015; Kuhlmann *et al.* 2016; Sammartini *et al.* 2018) have indicated a relationship between tephra layers or turbidites and slide failure planes. Although the exact nature of this relationship is disputed (Wiemer & Kopf 2016), it is thought that tephra could behave as “weak layers” – layers of inherently lower strength than adjacent layers, which are thereby prone to failure (Locat *et al.* 2014). Using core logging, sedimentological, and geotechnical data from the IODP Expedition 340, Lafuerza *et al.* (2014) showed that low hydraulic conductivity of hemipelagic sediments offshore Martinique (Lesser Antilles) could cause low rates of dewatering in turbidites and tephra layers, allowing excess pore fluid pressures to persist at depth. Moreover, Hornbach *et al.* (2015) suggested that even small changes in the stress regime of these layers, such as that resulting from regional strain and grain reorganization during the compaction of sediments, might trigger motion. Effectively, results from IODP 340 showed that sand layers (i.e. tephra ash and

turbidites), as well as boundaries between sand and mud layers, may act as multiple decollement surfaces that promote and enhance the mobility of landslide deposits (Le Friant *et al.* 2015).

### **Geological Setting**

The Cape Verdes archipelago, offshore West Africa (Figure 1), is the surface expression of a mantle hotspot (Crough 1978; Holm *et al.* 2008; Ramalho *et al.* 2010). The island of Fogo, in the southern part of the archipelago, is one of the most active oceanic intraplate volcanoes in the world, having erupted 28 times in the last 520 years (Ribeiro 1960; Torres *et al.* 1997; González *et al.* 2015). Such volcanically active areas are well known to be associated with seismicity and the southern Cape Verdes are no different, recording frequent volcano-tectonic earthquakes (Grevemeyer *et al.* 2010; Faria & Fonseca 2014; Vales *et al.* 2014). Sediment cores collected in the region provide evidence of at least 43 large, explosive eruptions in the area in the last 150 kyr (Eisele *et al.* 2015), attesting to Fogo's vigorous volcanic activity. Such frequent volcanic activity, along with the accumulation of volcanic deposits on the submarine flanks, means that Fogo's flanks are potentially unstable and prone to collapsing.

An up-to-1 km high, semi-circular depression (Bordeira), open to the east on central Fogo, was interpreted by Day *et al.* (1999) as the scar of a large flank-collapse, referred to as "Monte Amarelo" (Figure 2). Other authors, however, interpret the same morphology as two partially overlapping volcanic calderas that were later cut by a flank-collapse that affected the eastern portion of the edifice (Torres *et al.* 1997; Brum da Silveira *et al.* 1997; Madeira *et al.* 2008; Martínez-Moreno *et al.* 2018). Fogo's Monte Amarelo flank-collapse, however, is strongly supported by the presence of a preserved lateral ramp at Espigão (Brum da Silveira *et al.* 1997), and a debris avalanche deposit located between the islands of Fogo and Santiago (Le Bas *et al.* 2007; Masson *et al.* 2008) (Figures 2, 3). Using a combined magnetotelluric- and multibeam-based approach, Martínez-Moreno *et al.* (2018) estimated a volume of 110 km<sup>3</sup> for the Monte Amarelo debris avalanche deposit. This estimate corresponds well with previous bathymetric-based estimates, which ranged between 80 and 160 km<sup>3</sup> (Le Bas *et al.* 2007; Madeira *et al.* 2008; Masson *et al.* 2008). Tsunami deposits found on the nearby islands of Santiago and Maio indicate that the Monte Amarelo flank-collapse was tsunamigenic, with the resulting tsunami achieving a run-up in excess of 270 m above coeval sea level on Santiago (Ramalho *et al.* 2015; Madeira *et al.* 2019). The exact age of the collapse, however, is the topic of ongoing debate. On the basis of <sup>3</sup>He geochronology of lava flows from Fogo, Foeken *et al.* (2009) first estimated the collapse to have occurred between 62 and 123 ka. Paris *et al.* (2011) suggested a narrower window of 86 to 124 ka, based, respectively, on Ar/Ar ages of lava flows from Fogo thought to be post-collapse, and U-Th dating of corals from tsunami deposits on Santiago. More recently, based on the results of cosmogenic <sup>3</sup>He dating of tsunami megaclasts from Santiago Island, Ramalho *et al.* (2015) proposed that the collapse and ensuing tsunami took place between 65 and 84 ka, with a most probable age of 73 ± 7 ka. This agrees, within uncertainty, with the age recently reported by Maderia *et al.* (2019; 78 ± 0.9 ka) for a set of tsunami deposits found on the coast of Maio Island. Dating of turbidite material attributed to tsunami-triggered sediment transport along

the flanks of Fogo and Brava islands, however, led Eisele *et al.* (2015) to favour an older age of 86-117 ka. Finally, and most recently, Marques *et al.* (2019) suggested a much younger age of 43-59 ka, based on K/Ar dating of Fogo lava flows that they considered to be pre- and post-collapse.

Whether Fogo has collapsed only once or multiple times is also the subject of discussion (Day *et al.* 1999; Ramalho *et al.* 2015; Martínez-Moreno *et al.* 2018; Marques *et al.* 2019). Subsequent eruptions on Fogo have largely been constrained to the central and eastern parts of the island, and a prominent stratovolcano – Pico do Fogo – presently rises to 2829 m above sea level within the landslide scar (Figure 2; Torres *et al.* 1997).

### **Objectives**

A detailed analysis of the distribution of the landslide deposits and failure mechanism are critical for constraining the hazard linked to the collapse of a volcanic flank. Moreover, this characterization is crucial for tsunami hazard modelling, given that the mode, volume and run-out of a collapse will have a profound effect on the resulting tsunami waves (Grilli *et al.* 1997; Abadie *et al.* 2012; Watt *et al.* 2012a). The offshore distribution of the Monte Amarelo flank-collapse debris (i.e. debris avalanche deposits) and the surrounding volcanoclastic apron were previously mapped using only medium-resolution (100 m grid cell size) multibeam bathymetric data (Masson *et al.* 2008). Consequently, the distribution of the debris avalanche deposits, and of any additional slope instabilities in the area, are still poorly constrained. As Fogo's volcanic flank-collapse likely triggered a megatsunami with a recognizable impact on the adjacent islands' coastlines (Paris *et al.* 2011, 2018; Ramalho *et al.* 2015; Madeira *et al.* 2019), a full characterization of this landslide and its related volume is crucial for improving numerical models of tsunami generation, propagation and inundation, and for constraining the hazard potential associated with large, tsunamigenic volcanic flank-collapses.

Here, using recently acquired multibeam bathymetric data (50 m grid cell size) in conjunction with parametric sediment echo-sounder data and sediment gravity cores, we revisit the Monte Amarelo volcanic flank-collapse and consider general slope stability in the southern distal region of the volcanic apron. We aim to (i) map out the lateral extent and characterise the acoustic nature of the Monte Amarelo debris avalanche deposits; (ii) identify possible additional landslides on the slopes south of the islands of Fogo and Santiago; and (iii) consider the related preconditioning and triggering processes.

### **Data and methodology**

The data presented in this paper were collected during R/V Meteor cruise M155 (May - June 2019; Krastel *et al.* 2019) and are supplemented by multibeam bathymetric data collected during R/V Meteor cruise M80/3 (Hansteen *et al.* 2014). The bathymetric data from both cruises were acquired using hull-mounted Kongsberg EM120 (M80/3), EM122 (M155) and EM710 (M155) multibeam echo-sounders. The EM120/EM122 system has a swath coverage of up to 150° and a nominal sonar frequency of 12 kHz, and is designed to perform seabed mapping to full ocean depth. During cruise M155, the swath width was reduced to 120° in

order to increase the quality and resolution of the data. The EM710 system has a swath coverage of up to 140°, and a nominal sonar frequency of 70 to 100 kHz. As such, the EM710 was only used in water depths less than 700 m close to the islands of Fogo and Santiago. All bathymetric data were filtered for outliers and manually edited. The data from all cruises and multibeam systems are combined and gridded at 50 m.

Parametric sediment echo-sounder data were collected using a parametric ATLAS DS-3/P70 system (Parasound). This system has an opening angle of 4° and operates at primary high frequencies of 18.5 kHz and 22.5 kHz, resulting in a parametric low frequency of 4 kHz. The vertical resolution of this system is in the decimeter range. All depth scales on Parasound images presented in this study were calculated using a constant velocity of 1500 m/s.

Sediment cores were collected during cruise M155 using a gravity corer with tube lengths of 3 to 15 m. These cores were visually described onboard, and smear slides were analyzed to obtain further microscopic information about the core mineralogy, texture and composition. Dating and further geochemical and sedimentological analysis of these cores is ongoing and will form the basis of future work on the subject.

## Results

### *The Monte Amarelo debris avalanche deposits (MTD-A)*

The Monte Amarelo deposits proximal to Fogo (*MTD-A*; distribution shown in Figure 3a) are characterized by overlapping diffraction hyperbolae (Figure 3b). This hummocky character is typical of debris avalanche deposits from volcanic flank-collapses (Siebert 1984). The sedimentary drape covering *MTD-A* is relatively thin (<1.5 m thick) or not imaged, and is often characterized by two strong, positive reflections. Irregularly shaped blocks that are up to 100 m higher than the surrounding seafloor are present within the landslide debris (Figure 2). Further away from Fogo, the hummocky topography is less prevalent in the bathymetric data, despite the presence of diffraction hyperbolae in the Parasound data. This highlights a decrease in the size of the hummocks with distance from the island. We note a progressive transition in the acoustic signature of the deposits from a hyperbolic facies with metre-scale acoustic penetration (Figure 3b), to a mounded facies draped by ~1.5 m of sediment characterized by the aforementioned prominent double reflections (Figure 3d). In a few places, these double reflections are replaced by a succession of finely stratified layers (inset of Figure 3c).

### *Distal deposits related to the Monte Amarelo volcanic flank-collapse (MTD-B)*

Southwards, with increasing distance from Fogo, the acoustic character of the Monte Amarelo deposits changes from being hyperbolae-dominated to being characterised by an acoustically transparent/semi-transparent facies (Figure 3d; hereafter referred to as *MTD-B*; distribution shown in Figure 3a). In the northern reaches of *MTD-B*, the upper surface of the deposits is undulating, with metre-scale variations in depth over wavelengths of tens to hundreds of metres (Figure 3d). The overlying sedimentary drape mantles the surface of the deposits and a strong internal reflector is imaged in places within the deposits (Figure 3d).

This internal reflector clearly contrasts with the overlying transparent facies, separating it from the lower semi-transparent facies. In the northern and western reaches of *MTD-B*, this internal reflector is either clearly imaged or not present at all, and the base of the lower part of *MTD-B* is not resolved (Figure 3d).

In the eastern and southern reaches of *MTD-B*, the aforementioned internal reflector is continuous over a relatively long distance (Figure 3e). In these areas, a thicker sedimentary drape (up to ~4 m thick), containing multiple strong reflections alternating with relatively thin, transparent layers, overlies *MTD-B*. The upper surface of *MTD-B* is diffuse and undulating in parts. The base of *MTD-B* is marked by a prominent reflection that has a similar acoustic signature, although of lower amplitude, as the double reflector that overlies the deposits. A series of well-stratified sediments is imaged below the base of *MTD-B* (Figure 3e). At the eastern lateral margin of *MTD-B*, the internal reflector within the slide deposits is more diffuse (Figure 3e). The lowermost part of *MTD-B* remains transparent in nature, but remnant stratifications are imaged in some areas of the uppermost part. The eastern boundary is gradational over two to three km, and is marked by a progressive thinning of the transparent facies, and an increase in stratification within the deposits (Figure 3e).

Constraining the thickness and lateral extent of *MTD-B* is challenging. In the northern reaches, where the strong internal reflector is imaged, the base of the deposits is not resolved by the acoustic system (Figure 3c). This means that only the thickness of the upper transparent facies related to *MTD-B* (up to ~7 m thick, but variable) can be estimated. In its eastern and southern extents, however, the bases of the upper and lower units of *MTD-B* are imaged, leading to thickness estimates of 4-6 m and 5-15 m for the upper and lower units, respectively (Figure 3e). The large spacings between the Parasound profiles in the central and southern parts of the working area, south of Fogo and Santiago, mean that we cannot constrain the thickness of the deposits in these areas with certainty. In addition, *MTD-B* deposits might extend beyond the limits of the surveyed area, especially to the west and south (Figure 3a).

#### ***Additional mass wasting events on the shallow slopes south of Fogo and Santiago***

Visual analysis of gravity cores taken in the southern part of the volcanoclastic apron of Fogo and Santiago reveals mud- and nanofossil-rich facies interbedded with multiple sand units that are defined by parallel and cross-laminated, seldom normally-graded beds (Figure 4). These sandy intervals generally range from fine to coarse sand deposits, and are typically of centi- to decimetre scale (Figure 4b). Microscopic analyses show that these shallow sandy turbidites are predominantly of volcanoclastic (volcanic lithics that are mostly lava fragments, tachylitic to brown glass, crystal fragments), and/or mixed volcanoclastic-bioclastic composition, rather than solely bioclastic.

Multibeam-bathymetry and sediment echo-sounder data on the shallow slopes south of Fogo and Santiago reveal multiple morphological steps and several mass transport deposits (MTDs; Figure 5). These additional MTDs are characterized by acoustically transparent and/or hyperbolic facies and occur both above and below *MTD-B*. Several of these MTDs are

exposed at the surface and have remobilized previously failed sediments, including those of *MTD-B* (e.g. Figure 5, 6). However, as for *MTD-B*, their full lateral extent is generally poorly constrained due to large profile spacings and their possible continuation outside the surveyed area. The morphological steps vary in length, height, strike, and orientation (Figure 5a), but are typically near-vertical (Figure 5c, d). In the following section, we describe the character of, and relation between, several of these features in more detail.

#### *Tectonic escarpments along the western and eastern margins*

A series of escarpments is imaged to the west, along the western margin of *MTD-B*, and stretching between 40 and 110 km south of Fogo (Figure 5a, b). The most prominent escarpment is orientated roughly north-south and extends across ~55 km. It is ~50 m high and near vertical (Figure 5c). Monte Amarelo *MTD-B* deposits are imaged on the eastern side (footwall) of this step. On the western (hanging wall) side, the sediments are characterized by a semi-transparent facies with a hummocky upper surface (hereafter referred to as *MTD-X*; Figure 5c). In places, a pair of strong reflectors is identified within the sedimentary drape that overlies *MTD-X*. Proximal to the escarpment, the base of *MTD-X* cannot be resolved. Further west of the scarp, however, stratified sediment is imaged below the base of *MTD-X* (inset 1 of Figure 5c). The lateral extent of *MTD-X* is constrained to both the north and east by the morphological steps (Figure 5b, c), resulting in a minimum area of 900 km<sup>2</sup> for *MTD-X*.

#### *Southern Scour Complex*

A prominent, elongated scour-shaped feature is evident in the central part of the southern distal region (hereafter referred to as the *Southern Scour Complex (SSC)*, Figure 6a). This feature is ~60 km long and covers an area of ~340 km<sup>2</sup>. The western and eastern sides of the *SSC* are constrained by up-to-40 and 55 m high escarpments, respectively (Figure 6). In Parasound data crossing the complex, acoustically transparent facies with a strong internal reflector characterizing the Monte Amarelo *MTD-B* is evident on both the eastern and western sides of the *SSC* (Figure 6b, c). In the central part of the *SSC* (Figure 6c), acoustically transparent MTDs with some internal structure are covered by ~3 m of stratified sediment. In the northern part, the material within the *SSC* is characterized by a semi-transparent facies with a hummocky upper surface, and the deposits are largely exposed at the surface (Figure 6b). An additional instability is visible on the eastern side of the *SSC* (hereafter referred to as the *Eastern Scour (ES)*; Figure 6d). The *ES* reaches a thickness of up to 35 m thick in its centre, and covers an area of ~120 km<sup>2</sup>. Parasound data across the *ES* show that this failure primarily remobilized the Monte Amarelo *MTD-B* deposits in this region along the same basal glide plane as *MTD-B* (Figure 6d). To the south, the *SSC* is fan-shaped, with fingers of unfailed stratified sediments standing up to ~20 m above failed sediment (Figure 6d). Semi-transparent deposits characteristic of *MTD-B* are imaged on the tops of these stratified fingers.



## Discussion

### *The two-fold nature of Fogo's Monte Amarelo flank-collapse*

Based on the difference in their acoustic character, we distinguish between two main types of deposits related to the Monte Amarelo volcanic flank-collapse: hummocky, debris avalanche deposits proximal to Fogo (*MTD-A*), and finer-grained, acoustically transparent landslide deposits (*MTD-B*) at greater distances from Fogo that were previously not recognized (Figure 3). The blocky debris avalanche deposits of *MTD-A* cover a surface area of 3,180 km<sup>2</sup>. If we also include the region in which the size of the hummocks decreases with distance from the source (marked by the yellow shaded region in Figure 3a), this increases the total area of *MTD-A* to 6,820 km<sup>2</sup>; more than four times the previous estimate (~1,470 km<sup>2</sup>; Masson *et al.* 2008). As the base of the volcanic debris avalanche deposits is not resolved by the Parasound data, we cannot, at this stage, revisit estimates of the volume involved.

Monte Amarelo *MTD-B* is characterised by an acoustically transparent to semi-transparent facies that is overlain by a series of two to four strong reflectors (Figure 3, 5c, 6b-d). A prominent, internal reflection is clearly observed in places within the deposits; separating *MTD-B* into two main parts (Figure 3 d, e). The uppermost section is of relatively constant thickness (4-6 m). Contrastingly the thickness of the lower part of *MTD-B* is highly variable (up to 15 m thick). Interestingly, the upper and lower parts of *MTD-B* have a similar areal distribution within the volcanoclastic apron; covering a minimum of 18,400 km<sup>2</sup> (upper) and 19,500 km<sup>2</sup> (lower) (Figure 3a). This implies minimum volumes of 92 km<sup>3</sup> (upper) and 195 km<sup>3</sup> (lower), assuming average thicknesses of 5 m (upper) and 10 m (lower). Their similar distribution implies that the upper and lower parts of *MTD-B* might share the same source or, at least, be genetically linked. We therefore infer that the upper and lower part of *MTD-B* might have been emplaced (almost) contemporaneously. The absolute timing of these two phases, however, remains unclear. Of particular interest is whether these two failures occurred as two phases of the same event, or as two distinct events separated by some time. Further analytical work on the sediment cores will help to reveal more information about the nature of the prominent internal reflection, along with the relative timing of these two depositional episodes.

A similar two-fold nature of volcanic flank-collapse deposits – hummocky debris avalanche deposits accompanied by acoustically transparent to seismically chaotic deposits with a comparatively smooth upper surface – has also been reported for flank-collapses at other locations, including in the Lesser Antilles (Watt *et al.*, 2012a, b; Le Friant *et al.* 2015; Brunet *et al.* 2016); at La Réunion (Indian Ocean; Lebas *et al.* 2018); and at Ritter Island (Papua New Guinea; Karstens *et al.* 2019; Watt *et al.* 2019). Drilled cores retrieved during IODP Expedition 340 in the Lesser Antilles indicated that widespread, seismically chaotic deposits (interpreted as equivalent to our *MTD-B*) primarily consist of hemipelagic mud interbedded with a combination of tephra, volcanoclastic layers, or bioclastic turbiditic deposits, which have undergone varying degrees of deformation (Le Friant *et al.* 2015; Brunet *et al.* 2016). To explain these findings, Le Friant *et al.* (2015) proposed a failure model where the loading of seafloor sediment by volcanic debris avalanche deposits triggered sediment destabilization

and progressive downslope-propagating failure along a decollement. In this way, the deformation can propagate great distances away from the flank collapse, affecting seafloor sediments that were otherwise stable (Le Friant *et al.* 2015). The acoustic character of our *MTD-B* bears a strong resemblance to the seismically chaotic deposits in the Lesser Antilles. We, therefore, interpret *MTD-B* to be the result of the failure of pre-existing seafloor sediments following the loading of the Monte Amarelo debris avalanche deposits (*MTD-A*).

### ***A history of mass-wasting and remobilisation in the southern Cape Verdes***

#### *Repeated mass-wasting events at Fogo?*

Preliminary stratigraphic correlations based on visual similarities of cores in the southern distal part of the working area indicate at least seven volcanoclastic or mixed volcanoclastic-bioclastic sandy turbidite layers above and below the Monte Amarelo flank-collapse deposits (Figure 4). These turbidite layers are dominated by mafic glass, crystals and lava fragments, which, based on their petrography, suggests that they originated from Fogo volcano. This indicates that smaller mass-wasting events may have occurred at Fogo in addition to the Monte Amarelo volcanic flank-collapse, as also recently suggested by Marques *et al.* (2019). Correlation and origin of these volcanoclastic deposits will, however, be verified by future analytical work on the sediment cores.

#### *Regional tectonic and volcanic influences south of Fogo?*

The multiple fault-related escarpments and MTDs identified in the acoustic data south of Fogo further highlight a long and complex history of instability and slope failure in the region (Figure 5). Ramalho *et al.* (2010) found that the island of Santiago has uplifted at a rate of ~100 m/Myr over the past 4 Myr, and attributed this uplift to a combination of magmatic intrusions under or within the island edifice, together with episodic swell-wide uplift that affected the whole archipelago. Although geologically recent uplift has not been reported at Fogo, the neighbouring island of Brava has experienced one of the most dramatic intrusion-related uplift trends of any ocean island in the world, with up to 400 m of uplift in the last 1.8 Ma (Madeira *et al.* 2010). The presence of such widespread uplift across the archipelago – and particularly around the southern Cape Verdes – thus suggests that the series of escarpments and landslides observed on the slopes south of Fogo may be the surface expression of a combination of regional tectonics, associated with the growth of the Cape Verdes hotspot swell, and crustal intrusions in the vicinity of the islands. As several of the exposed faults have landslide deposits on their hanging wall side, we interpret that these MTDs are most likely the results of movement along the faults and, as such, the result of neo-tectonic activity.

The *Southern Scour Complex (SSC)*, which is exposed at the seafloor, post-dates the Monte Amarelo flank-collapse (Figure 6). Monte Amarelo *MTD-B* deposits are imaged on the footwall east of the *SSC* (Figure 6b-d), and on the tops of the unfailed, stratified fingers at the southern extent of the *SSC* (Figure 6d). This indicates that the *SSC* formed after the Monte Amarelo flank-collapse, remobilizing *MTD-B*. In the northern and eastern parts of the *SSC*, acoustically transparent mass-wasting deposits are exposed at the seafloor, highlighting more

recent mass wasting and remobilization of *MTD-B* (Figure 6b). We interpret the more recent mass-wasting deposits in the northern and eastern parts of the *SSC* as evidence of retrogressive development of the *SSC* in these directions (Figure 6e).

The presence of fault-related escarpments on the slopes south of Fogo, frequent volcanic episodes at Fogo (Ribeiro 1960; Torres *et al.* 1997), and ongoing uplift at the neighbouring islands of Santiago and Brava (Madeira *et al.* 2010; Ramalho *et al.* 2010), together imply the occurrence of frequent earthquakes in the past. The volcanic activity at Fogo has resulted in the widespread deposition of numerous tephra layers (Eisele *et al.* 2015). Subsequent earthquake shaking may have resulted in liquefaction of overlying layers, which, together with sedimentary over-pressure following further sediment deposition, can help to precondition the shallow slopes for failure (Moernaut *et al.* 2019). Even small changes in the stress regime of tephra layers have been found to be able to trigger failure (Hornbach *et al.* 2015). Consequently, it is possible that some of the MTDs observed on the shallowly dipping slopes south of Fogo, and unrelated to escarpments, may result from such a process. Future work, particularly on the sediment cores collected during cruise M155, will help to shed light on the sources and processes responsible for the MTDs mapped on the shallow slopes south of Fogo and Santiago.

### **Implications and Conclusions**

In this study, we revisited the Monte Amarelo volcanic flank-collapse of Fogo Island and found that the deposition of the debris avalanche material may have triggered subsequent failures of pre-existing seafloor sediments in (at least) two phases. This is similar to what has been observed in the Lesser Antilles (Le Friant *et al.* 2015; Brunet *et al.* 2016), at La Réunion (Lebas *et al.* 2018), and at Ritter Island (Karstens *et al.* 2019; Watt *et al.* 2019). It is not yet clear whether these two phases of seafloor sediment failure were synchronous as a result of a single flank-collapse, or if they reflect multiple, distinct events. The question of whether the main flank-collapse occurred as a single or as multiple events is of utmost importance for tsunami modelling because the volume and timing of individual failures are the main factors controlling the tsunamigenic potential (Løvholt *et al.* 2015). Water depth and landslide kinematics play a key role in controlling the tsunamigenic potential of a subaqueous landslide (Watts *et al.* 2000; Ward, 2001; Watt *et al.* 2012a; Harbitz *et al.* 2014). As *MTD-B* occurred in depths exceeding 3,000 m, we consider that its influence on the resulting tsunami magnitude was negligible, and that only the hummocky debris avalanche deposits (*MTD-A*) contributed to the megatsunami that inundated nearby islands.

Multibeam bathymetric, sediment echo-sounder, and sediment gravity core data acquired during cruise M155 of R/V Meteor provide seafloor evidence that show, for the first time, the presence of multiple additional mass-wasting events on the shallow slopes south of Fogo. Turbidite sands recovered in the cores have a volcanoclastic or mixed volcanoclastic-bioclastic composition, which suggests that prevailing mass-transport processes in the region may be dominated by volcanic eruptions or volcanic mass-wasting events. However, it is also possible that some of the additional mass-wasting events are unrelated to flank-collapse events or submarine failures, and could instead have been generated by sedimentary

overpressure and/or by the presence of a mechanically weak layer such as tephra or turbidites. Further analytical work is required to verify the sources of these events on the shallow slopes south of the Cape Verdean archipelago. As these layers are thin, have limited lateral extent, and occurred on low-gradient slopes in deep water, we consider their associated tsunamigenic potential to be low.

### **Acknowledgements**

We are grateful to the captain, crew and scientific teams of R/V Meteor (M155) for their enthusiasm and assistance onboard. We are also thankful for the thoughtful reviews of the editor, Lawrence Amy, as well as Paraskevi Nomikou and two anonymous reviewers, whose comments improved this manuscript.

### **Funding information**

Data acquisition during R/V Meteor's cruise M155 was funded by the DFG and BMBF. R. Barrett is funded by the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 721403 (SLATE – “Submarine Landslides and Their Impact on European Continental Margins”). R. Ramalho acknowledges his IF/01641/2015 contract funded by Fundação para a Ciência e Tecnologia (FCT), as well as the following projects funded by FCT: IF/01641/2015, LISBOA-01-0145-FEDER-028588 and PTDC/CTA-GEO/28588/2017 UNTIeD, PTDC/GEO-GEO/1123/2014 FIRE, and UID/GEO/50019/2019 - Instituto Dom Luiz.

**Correction notice:** The original Accepted Manuscript posted on 19 December 2019 was incorrect. This was due to a publisher error resulting in processing a version of the manuscript that was not finalized by the authors.

### **References**

- Abadie, S.M., Harris, J.C., Grilli, S.T., & Fabre, R., 2012, Numerical modeling of tsunami waves generated by the flank collapse of the Cumbre Vieja Volcano (La Palma, Canary Islands): tsunami source and near field effects. *Journal of Geophysical Research: Oceans*, v. 117, n. C05030, p. 1-26, <https://doi.org/10.1029/2011JC007646>.
- Begét, J. E., & Kienle, J., 1992, Cyclic formation of debris avalanches at Mount St Augustine volcano. *Nature*, v. 356, p. 701-704, <https://doi.org/10.1038/356701a0>.
- Blahůt, J., Balek, J., Klimeš, J., Rowberry, M., Kusák, M., & Kalina, J., 2019, A comprehensive global database of giant landslides on volcanic islands, *Landslides*, v. 16, n. 10, p. 2045-2052, <https://doi.org/10.1007/s10346-019-01275-8>.
- Brum da Silveira, A., Madeira, J., Serralheiro, A., Torres, P.C., da Silva, L.C., Mendes, M.H., 1997, O controlo estrutural da erupção de Abril de 1995 na Ilha do Fogo, Cabo Verde. *In A Erupção Vulcânica de 1995 na Ilha do Fogo, Cabo Verde* (Publ. IICT, Lisboa, Portugal), p. 51–61.

Brunet, M., Le Friant, A., Boudon, G., Lafuerza, S., Talling, P., Hornbach, M., Ishizuka, O., Lebas, E., Guyard, H., & IODP Expedition 340 Science Party, 2016, Composition, geometry, and emplacement dynamics of a large volcanic island landslide offshore Martinique: From volcano flank- collapse to seafloor sediment failure? *Geochem. Geophys. Geosyst.*, v. 17, n. 3, p. 699-724, <https://doi.org/10.1002/2015GC006034>.

Casalbore, D., Romagnoli, C., Bosman, A., & Chiocci, F. L., 2011, Potential tsunamigenic landslides at Stromboli Volcano (Italy): Insight from marine DEM analysis. *Geomorphology*, v. 126, n. 1-2, p. 42-50, <https://doi.org/10.1016/j.geomorph.2010.10.026>.

Casalbore, D., Romagnoli, C., Pimentel, A., Quartau, R., Casas, D., Ercilla, G., Hipólito, A., Sposato, A., & Chiocci, F. L., 2015, Volcanic, tectonic and mass-wasting processes offshore Terceira Island (Azores) revealed by high-resolution seafloor mapping. *Bulletin of Volcanology*, v. 77, n. 24, p. 1-19, <https://doi.org/10.1007/s00445-015-0905-3>.

Casalbore, D., 2018, Volcanic islands and seamounts. *In* *Submarine Geomorphology*, Springer, Cham, p. 333-347, [https://doi.org/10.1007/978-3-319-57852-1\\_17](https://doi.org/10.1007/978-3-319-57852-1_17).

Cervelli, P., Segall, P., Johnson, K., Lisowski, M., & Miklius, A., 2002, Sudden aseismic fault slip on the south flank of Kilauea volcano. *Nature*, v. 415, p. 1014-1018, <https://doi.org/10.1038/4151014a>.

Clare, M. A., Le Bas, T., Price, D., Hunt, J., Sear, D., Cartigny, M. J. B., Vellinga, A., Symons, W., Firth, C., & Cronin, S. J., 2018, Complex and cascading triggering of submarine landslides and turbidity currents at volcanic islands revealed from integration of high-resolution onshore and offshore surveys. *Frontiers in Earth Science*, v. 6, n. 223, <https://doi.org/10.3389/feart.2018.00223>.

Crough, S.T., 1978, Thermal origin of mid-plate hot-spot swells. *Geophys. J. R. astr. Soc.*, v. 55, p. 451-469, <https://doi.org/10.1111/j.1365-246X.1978.tb04282>.

Day, S. J., Heleno da Silva, S. I. N., & Fonseca, J. F. B. D., 1999, A past giant lateral collapse and present-day flank instability of Fogo, Cape Verde Islands. *Journal of Volcanology and Geothermal Research*, v. 94, p. 191-218, [https://doi.org/10.1016/S0377-0273\(99\)00103-1](https://doi.org/10.1016/S0377-0273(99)00103-1).

Eisele, S., Reißig, S., Freundt, A., Kutterolf, S., Nürnberg, D., Wang, K. L., & Kwasnitschka, T., 2015, Pleistocene to Holocene offshore tephrostratigraphy of highly explosive eruptions from the southwestern Cape Verde Archipelago, *Marine Geology*, v. 369, p. 233-250, <http://dx.doi.org/10.1016/j.margeo.2015.09.006>.

Faria, B., & Fonseca, J. F. B. D., 2014, Investigating volcanic hazard in Cape Verde Islands through geophysical monitoring: network description and first results. *Nat. Hazards Earth Syst. Sci.*, v. 14, p. 485-499, <https://doi.org/10.5194/nhess-14-485-2014>.

Foeken, J. P. T., Day, S., & Stuart, F. M., 2009, Cosmogenic <sup>3</sup>He exposure dating of the Quaternary basalts from Fogo, Cape Verdes: Implications for rift zone and magmatic

reorganization, *Quaternary Geochronology*, v. 4, n. 1, p. 37–49, <https://doi.org/10.1016/j.quageo.2008.07.002>.

Goff, J., Terry, J. P., Chagué-Goff, C., & Goto, K., 2014, What is a mega-tsunami? *Marine Geology*, v. 358, p. 12-17, <https://doi.org/10.1016/j.margeo.2014.03.013>.

González, P.J., Bagnardi, M., Hooper, A.J., Larsen, Y., Marinkovic, P., Samsonov, S.V., & Wright, T.J., 2015, The 2015-2015 eruption of Fogo volcano: Geodetic modeling of Sentinel-1 TOPS interferometry. *Geophysical Research Letters*, v. 42, n. 21, p. 9239-9246, <https://doi.org/10.1002/2015GL066003>.

Grevemeyer, I., Helffrich, G., Faria, B., Booth-Rea, G., Schnabel, M., & Weinrebe, W., 2010, Seismic activity at Cadamosto seamount near Fogo Island, Cape Verdes – formation of a new ocean island? *Geophysical Journal International*, v. 180, n. 2, p. 552-558, <https://doi.org/10.1111/j.1365-246X.2009.04440.x>.

Grilli, S.T., Svendsen, I.A. and Subramanya, R., 1997, Breaking criterion and characteristics for solitary waves on slopes. *Journal of waterway, port, coastal, and ocean engineering*, v. 123, n. 3, p.102-112, [https://doi.org/10.1061/\(ASCE\)0733-950X\(1997\)123:3\(102\)](https://doi.org/10.1061/(ASCE)0733-950X(1997)123:3(102)).

Grilli, S. T., Tappin, D. R., Carey, S., Watt, S. F. L., Ward, S. N., Grilli, A. R., Engwell, S. L., Zhang, C., Kirby, J. T., Schambach, L., & Muin, M., 2019, Modelling of the tsunami from the December 22, 2018 lateral collapse of Anak Krakatau volcano in the Sunda Straits, Indonesia. *Sci. Rep.*, 9, 11946, <https://doi.org/10.1038/s41598-019-48327-6>.

Gross, F., Krastel, S., Chiocci, F. L., Ridente, D., Bialas, J., Schwab, J., Beier, J., Cukur, D., & Winkelmann, D., 2014, Evidence for Submarine Landslides offshore Mt. Etna, Italy. *In* Krastel *et al.* (eds) *Submarine Mass Movements and Their Consequences. Advances in Natural and Technological Hazards Research*, Springer, Cham, v. 37, p. 307-316, [https://doi.org/10.1007/978-3-319-00972-8\\_27](https://doi.org/10.1007/978-3-319-00972-8_27).

Hansteen, T. H., Kwasnitschka, T., & Klügel, A., 2014, Cape Verde Seamounts – Cruise No. M80/3 – December 29, 2009 – February 1, 2010 - Dakar (Senegal) - Las Palmas de Gran Canaria (Spain). *METEOR-Berichte*, M80/3, 42 pp., DFG-Senatskommission für Ozeanographie, [https://doi.org/10.2312/cr\\_m80\\_3](https://doi.org/10.2312/cr_m80_3).

Harbitz, C. B., Løvholt, F., & Bungum, H., 2014, Submarine landslide tsunamis: How extreme and how likely? *Natural Hazards*, v. 72, p. 1341-1374, <https://doi.org/10.1007/s11069-013-0681-3>.

Harders, R., Kutterolf, S., Hensen, C., Moerz, T., & Brueckmann, W., 2010, Tephra layers: A controlling factor on submarine translational sliding? *Geochemistry, Geophysics, Geosystems*, v. 11, n. 5, p. 1-18, <https://doi.org/10.1029/2009GC002844>.

Holm, P. M., Grandvuinet, T., Friis, J., Wilson, J. R., Barker, A. K., & Plesner, S., 2008, An  $^{40}\text{Ar}$ - $^{39}\text{Ar}$  study of the Cape Verde hot spot: Temporal evolution in a semistationary plate

environment, *J. Geophys. Res.*, v. 113, n. B08201, p. 1-22,  
<https://doi.org/10.1029/2007JB005339>.

Hornbach, M. J., Manga, M., Genecov, M. *et al.* 2015, Permeability and pressure measurements in Lesser Antilles submarine slides: Evidence for pressure-driven slow-slip failure. *J. Geophys. Res. Solid Earth*, v. 120, n. 12, p. 7986–8011,  
<https://doi.org/10.1002/2015JB012061>.

Karstens, J., Berndt, C., Urlaub, M., Watt, S. F. L., Micallef, A., Ray, M., Klaucke, I., Muff, S., Klaeschen, D., Kühn, M., Roth, T., Böttner, C., Schramm, B., Elger, J., & Brune, S., 2019, From gradual spreading to catastrophic collapse – Reconstruction of the 1888 Ritter Island volcanic sector collapse from high-resolution 3D seismic data. *Earth and Planetary Science Letters*, v. 517, p. 1-13, <https://doi.org/10.1016/j.epsl.2019.04.009>.

Krastel, S., Schmincke, H.-U., Jacobs, C. L., Rihm, R., Le Bas, T. P., & Alibes, B., 2001, Submarine landslides around the Canary Islands, *J. Geophys. Res.*, v. 106, p. 3977-3998,  
<https://doi.org/10.1029/2000JB900413>.

Krastel, S., Andrade, M., Barrett, R. *et al.*, 2019, The tsunamigenic gravitational flank-collapse of Fogo volcano, Cape Verde Islands, Seismic pre-site survey for an IODP site on the Cape Verde Plateau, Cruise No. M155, 26.05.19 - 30.06.19, Pointe-à-Pitre (Guadeloupe) - Mindelo (Cape Verde). METEOR-Berichte, Gutachterpanel Forschungsschiffe, Bonn, v. M155, p. 1-44, [https://doi.org/10.2312/cr\\_m155](https://doi.org/10.2312/cr_m155).

Kuhlmann, J., Huhn, K., & Ikari, M.J., 2016, Do embedded volcanoclastic layers serve as potential glide planes? An integrated analysis from the Gela Basin offshore southern Sicily. *In* Lamarche, G. *et al.* (Eds.), *Submarine Mass Movements and Their Consequences. Advances in Natural and Technological Hazards Research*, v. 41, Springer International Publishing, Cham, p. 273–280, [https://doi.org/10.1007/978-3-319-20979-1\\_27](https://doi.org/10.1007/978-3-319-20979-1_27).

Lafuerza S., Le Friant, A., Manga, M., Boudon, G., Villemant, B., Stroncik, N., Voight, B., Hornbach, M., Ishizuka, O., & the Expedition 340 Scientific Party, 2014, Geomechanical characterizations of submarine volcano flank sediments, Martinique, Lesser Antilles Arc. *In* Krastel, S., *et al.* (Eds.), *Submarine mass movements and consequences, Advances in Natural and Technological Hazards Research*, Springer International Publishing, Switzerland, v. 2014, n. 37, [https://doi.org/10.1007/978-3-319-00972-8\\_7](https://doi.org/10.1007/978-3-319-00972-8_7).

Lebas, E., Le Friant, A., Boudon, G., Watt, S.F.L., Talling, P.J., Feuillet, N., Deplus, C., Berndt, C., & Vardy, M.E., 2011, Multiple widespread landslides during the long-term evolution of a volcanic island: Insights from high-resolution seismic data, Montserrat, Lesser Antilles. *Geochemistry, Geophysics, Geosystems*, v. 12, n. 5,  
<https://doi.org/10.1029/2010GC003451>.

Lebas, E., Le Friant, A., Deplus, C., & de Voogd, B., 2018, Understanding the evolution of an oceanic intraplate volcano from seismic reflection data: A new model for La Réunion, Indian Ocean. *J. Geophys. Res.*, v. 123, p. 1035-1059, <https://doi.org/10.1002/2017JB014959>.

Le Bas, T.P., Masson, D.G., Holtom, R.T., & Grevemeyer, I., 2007, Slope failures of the flanks of the southern Cape Verde Islands. *In* Lykousis, V., Sakellariou, D., Locat, J. (eds.) *Submarine Mass Movements and Their Consequences. Advances in Natural and Technological Hazards Research*, Springer, Dordrecht, v. 27, p. 337-345, <https://doi.org/10.1007/978-1-4020>.

Le Friant, A., Ishizuka, O., Boudon, G. *et al.*, 2015, Submarine record of volcanic island construction and collapse in the Lesser Antilles arc: First scientific drilling of submarine volcanic island landslides by IODP Expedition 340. *Geochemistry, Geophysics, Geosystems*, v. 16, p. 420-442, <http://doi.org/10.1002/2014GC005652>.

León, R., Somoza, L., Urgeles, R. *et al.*, 2017, Multi-event oceanic island landslides: New onshore-offshore insights from El Hierro Island, Canary Archipelago. *Marine Geology*, v. 393, p. 156-175, <http://dx.doi.org/10.1016/j.margeo.2016.07.001>.

Livanos I., Nomikou P., Papanikolaou D., & Rousakis G., 2013. The volcanic debris avalanche on the SE submarine slope of Nisyros volcano, Greece: Geophysical exploration and implications for subaerial eruption history. *Geo-Marine Letters*, v. 33, n. 6, p. 419, <https://doi.org/10.1007/s00367-013-0338-y>.

Locat, J., Leroueil, S., Locat, A., & Lee, H. J., 2014, Weak Layers: Their Definition and Classification from a Geotechnical Perspective. *In* Krastel *et al.* (eds) *Submarine Mass Movements and Their Consequences. Advances in Natural and Technological Hazards Research*, Springer, Cham, v. 37, p. 3-12, [https://doi.org/10.1007/978-3-319-00972-8\\_1](https://doi.org/10.1007/978-3-319-00972-8_1).

Løvholt, F., Pedersen, G., Harbitz, C. B., Glimsdal, S., Kim, J., 2015, On the characteristics of landslide tsunamis. *Philosophical Transactions of the Royal Society A*, v. 373, n. 20140376, <http://dx.doi.org/10.1098/rsta.2014.0376>.

Madeira, J., Brum da Silveira, A., Mata, J., Mourão, C., & Martins, S., 2008, The role of mass movements on the geomorphologic evolution of island volcanoes: examples from Fogo and Brava in the Cape Verde archipelago. *Comunicações Geológicas*, v. 95, p. 93–106.

Madeira, J., Mata, J., Mourão, C., Brum dal Silveira, A., Martins, S., Ramalho, R., Hoffman, D.L., 2010, Volcano-stratigraphic and structural evolution of Brava Island (Cape Verde) based on  $^{40}\text{Ar}/^{39}\text{Ar}$ , U–Th and field constraints. *Journal of Volcanology and Geothermal Research*, v. 196, p. 219–235, <https://doi.org/10.1016/j.jvolgeores.2010.07.010>.

Madeira, J., Ramalho, R. S., Hoffmann, D. L., Mata, J., & Moreira, M., 2019, A geological record of multiple Pleistocene tsunami inundations in an oceanic island: The case of Maio, Cape Verde, *Sedimentology*, <https://doi.org/10.1111/sed.12612>.

Marques, F.O., Hildenbrand, A., Victória, S.S., Cunha, C., & Dias, P., 2019, Caldera or flank collapse in the Fogo volcano? What age? Consequences for risk assessment in volcanic islands. *Journal of Volcanology and Geothermal Research*, in press, <https://doi.org/10.1016/j.jvolgeores.2019.106686>.



Martínez-Moreno, F. J., Monteiro Santos, F. A., Madeira, J., Pous, J., Bernardo, I., Soares, A., Esteves, M., Adão, F., Ribeiro, A., Mata, J., & Brum da Silveira, A., 2018, Investigating collapse structures in oceanic islands using magnetotelluric surveys: The case of Fogo Island in Cape Verde, *J. Volc. and Geotherm. Res.*, v. 357, p.152-162, <https://doi.org/10.1016/j.jvolgeores.2018.04.028>.

Masson, D.G., Watts, A. B., Gee, M. J. R., Urgeles, R., Mitchell, N. C., Le Bas, T. P., & Canals, M. 2002, Slope failures on the flanks of the western Canary Islands. *Earth-Sci Rev*, v. 57, p. 1-35.

Masson, D. G., Le Bas, T. P., Grevemeyer, I., & Weinrebe, W., 2008, Flank collapse and large-scale landsliding in the Cape Verde Islands, off West Africa. *Geochemistry, Geophysics, Geosystems*, v. 9, n. 7, pp. 16, <https://doi.org/10.1029/2008GC001983>.

McGuire, W. J., 1996, Volcano instability: a review of contemporary themes. Geological Society, London, Special Publications, v. 110, p. 1-23, <https://doi.org/10.1144/GSL.SP.1996.110.01.01>.

McGuire, W. J., 2003, Volcano instability and lateral collapse. *Revista*, v. 1, p. 33-45.

McMurtry, G. M., Fryer, G. J., Tappin, D. R., Wilkinson, I. P., Williams, M., Fietzke, J., Garbe-Schoenberg, D., & Watts, P., 2004, Megatsunami deposits on Kohala volcano, Hawaii, from flank collapse of Mauna Loa. *Geology*, v. 32, p. 741-744, <https://doi.org/10.1130/G20642.1>.

Moernaut, J., Van Daele, M., Heirman, K., Wiemer, G., Molenaar, A., Vandorpe, T., Melnick, D., Hajdas, I., Pino, M., Urrutia, R., & De Batist, M., 2019, The subaqueous landslide cycle in south-central Chilean lakes: The role of tephra, slope gradient and repeated seismic shaking. *Sedimentary Geology*, v. 381, p. 84-105, <https://doi.org/10.1016/j.sedgeo.2019.01.002>.

Moore, J. G., Clague, D. A., Holcomb, R. T., Lipman, P. W., Normark, W. R., & Torresan, M. E., 1989, Prodigious submarine landslides on the Hawaiian Ridge. *J. Geophys. Res.*, v. 94, n. B12, p. 17,465-17,485, <https://doi.org/10.1029/JB094iB12p17465>.

Moore, J. G., & Moore, G. W., 1984, Deposit from a giant wave on the island of Lanai, Hawaii. *Science*, v. 226, n. 4680, p. 1312-1315, <https://doi.org/10.1126/science.226.4680.1312>.

Murray, J. B., & Voight, B., 1996, Slope stability and eruption prediction on the eastern flank of Mount Etna. Geological Society, London, Special Publications v. 110, p. 111-114, <https://doi.org/10.1144/GSL.SP.1996.110.01.08>.

Paris, R., Geichetti, T., Chevalier, J., Guillou, H., & Frank, N., 2011, Tsunami deposits in Santiago Island (Cape Verde archipelago) as possible evidence of a massive flank failure of Fogos volcano, *Sedimentary Geology*, v. 239, p. 129-145, <https://doi.org/10.1016/j.sedgeo.2011.06.006>.

Paris, R., Ramalho, R. S., Madeira, J., Ávila, S., May, S. M., Rixhon, G., Engel, M., Brückner, H., Herzog, M., Schukraft, G., Perez-Torrado, F. J., Rodriguez-Gonzalez, A., Carracedo, J. C., & Giachetti, T., 2018, Mega-tsunami conglomerates and flank collapses of ocean island volcanoes, *Marine Geology*, v. 395, p. 168-187, <http://dx.doi.org/10.1016/j.margeo.2017.10.004>.

Ramalho, R., Helffrich, G., Cosca, M., Vance, D., Hoffmann, D., & Schmidt, D. N., 2010, Episodic swell growth inferred from variable uplift of the Cape Verde hotspot islands. *Nature Geoscience*, v. 3, p. 774-777, <https://doi.org/10.1038/NGEO982>.

Ramalho, R. S., Winckler, G., Madeira, J., Helffrich, G. R., Hipólito, A., Quartau, R., Adena, K., & Schaefer, J. M., 2015, Hazard potential of volcanic flank collapses raised by new megatsunami evidence, *Scientific Advances*, v. 1, n. 9, e1500456, <https://doi.org/10.1126/sciadv.1500456>.

Ribeiro, O., 1960, *A ilha do Fogo e as suas erupções*. 2nd edn. Memórias, série geográfica I. Publ. Junta de Investigações do Ultramar, Ministério do Ultramar, Lisbon, Portugal.

Sammartini, M., Camerlenghi, A., Budillon, F., Insinga, D. D., Zgur, F., Conforti, A., Iorio, M., Romeo, R., & Tonielli, R., (2018), Open-slope, translational submarine landslide in a tectonically active volcanic continental margin (Licosa submarine landslide, southern Tyrrhenian Sea). *Geological Society, London, Special Publications*, v. 477, <https://doi.org/10.1144/SP477.34>.

Siebert, L., 1984, Large volcanic debris avalanches: characteristics of source areas, deposits, and associated eruptions. *Journal of Volcanology and Geothermal Research*, v. 22, p. 163-197, [https://doi.org/10.1016/0377-0273\(84\)90002-7](https://doi.org/10.1016/0377-0273(84)90002-7).

Tibaldi, A., 2001, Multiple sector collapses at Stromboli volcano, Italy: How they work. *Bulletin of Volcanology*, v. 63, p. 112-125, [https://doi.org/10.1016/S1464-1909\(99\)00022-2](https://doi.org/10.1016/S1464-1909(99)00022-2).

Tibaldi, A., Pasquarè, F.A., Papanikolaou, D., & Nomikou, P., 2008, Discovery of a huge sector collapse at the Nisyros volcano, Greece, by on-land and offshore geological-structural data. *Journal of Volcanology and Geothermal Research*, v. 177, n. 2, p. 485, <https://doi.org/10.1016/j.jvolgeores.2008.06.014>.

Torres, P., Madeira, J., Silva, L., Brum da Silveira, A., Serralheiro, A., & Mota Gomes, A., 1997, Carta geológica das erupções históricas da Ilha do Fogo: revisão e actualização: A erupção vulcânica de 1995 na Ilha do Fogo, Cabo Verde. *Publ IICT, Lisboa*, p. 119–132.

Vales, D., Dias, N. A., Rio, I., Matias, L., Silveira, G., Madeira, J., Weber, M., Carrilho, F., & Haberland, C., 2014, Intraplate seismicity across the Cape Verde swell: A contribution from a temporary seismic network. *Tectonophysics*, v. 636, p. 325-337, <https://doi.org/10.1016/j.tecto.2014.09.014>.

Ward S. N., 2001, Landslide tsunami. *Journal of Geophysical Research*, v. 106, n. B6, p. 11201–11215, <https://doi.org/10.1029/2000JB900450>.

Watt, S. F. L., Talling, P.J., *et al.* 2012a, Combinations of volcanic-flank and seafloor-sediment failure offshore Montserrat, and their implications for tsunami generation. *Earth Planet. Sci. Lett.*, v. 319, n. 320, p. 228–240, <https://doi.org/10.1016/j.epsl.2011.11.032>.

Watt, S. F. L., Talling, P.J., *et al.* 2012b, Widespread and progressive seafloor-sediment failure following volcanic debris avalanche emplacement: Landslide dynamics and timing offshore Montserrat, Lesser Antilles. *Marine Geology*, v. 323-325, p. 69-94, <https://doi.org/10.1016/j.margeo.2012.08.002>.

Watt, S. F. L., Karstens, J., Micallef, A., Berndt, C., Urlaub, M., Ray, M., Desai, A., Sammartini, M., Klauke, I., Böttner, C., Day, S., Downes, H., Kühn, M., & Elger, J., 2019, From catastrophic collapse to multi-phase deposition: Flow transformation, seafloor interaction and triggered eruption following a volcanic-island landslide. *Earth and Planetary Science Letters*, v. 517, p. 135-147, <https://doi.org/10.1016/j.epsl.2019.04.024>.

Watts, P., Imamura, F., & Grilli, S. T., 2000, Comparing model simulations of three benchmark tsunami generation cases. *Science of Tsunami Hazards*, v. 18, n. 2, p. 107–124.

Wiemer, G., & Kopf, A., 2016, On the role of volcanic ash deposits as preferential submarine slope failure planes. *Landslides*, v. 14, n. 1, <https://doi.org/10.1007/s10346-016-0706-6>.

Wiemer, G., Moernaut, J., Stark, N., Kempf, P., De Batist, M., Pino, M., Urrutia, R., Strasser, M., & Kopf, A., 2015, The role of sediment composition and behaviour under dynamic loading conditions on slope failure initiation: a study of a subaqueous landslide in earthquake-prone South-Central Chile. *International Journal of Earth Sciences*, v. 104, n. 5, p. 1439-1457, <https://doi.org/10.1007/s00531-015-1144-8>.

## Figure captions

**Fig. 1.** (a) Location of the Cape Verdes offshore West Africa. Box outlines the region presented in b. (b) Multibeam bathymetric data collected during R/V Meteor cruises M155 and M80/3 overlain on GEBCO2019 (British Oceanographic Data Centre, [www.gebco.net](http://www.gebco.net)) bathymetry of the southern Cape Verdean archipelago. Black lines show the locations of Parasound profiles collected during cruise M155 of R/V Meteor. (c) Location of the newly acquired Parasound data and of the gravity cores collected during cruise M155.

**Fig. 2.** High-resolution, multibeam bathymetry data highlighting the hummocky nature of the Monte Amarelo debris avalanche deposits (*MTD-A*) proximal to Fogo. (a) Compilation of the M155 and M80/3 multibeam bathymetry data overlain on GEBCO2019 bathymetry and DEM topography of Fogo Island. (1) Espigão escarpment; (2) Bordeira escarpment (dotted black line) is regarded as either the Monte Amarelo headwall scarp (Day *et al.* 1999), or the edge of a caldera (Madeira *et al.* 2008; Martínez-Moreno *et al.* 2018); (3) Monte Amarelo headwall scarp (dotted pink line; Madeira *et al.* 2008; Martínez-Moreno *et al.* 2018); dashed black line: previously estimated extent of the Monte Amarelo debris avalanche deposits (Masson *et al.* 2008); solid white line: revised extent of the Monte Amarelo debris avalanche deposits (*MTD-A*); dashed white line: transitional zone where the size of the hummocks decreases with distance from the source. (b) Enlarged view of the hummocky topography of the Monte Amarelo debris avalanche deposits. Note the angular shape and variable size of the hummocks.

**Fig. 3.** (a) Map showing the distribution of the debris avalanche deposits (*MTD-A*), the acoustically transparent/seismically chaotic *MTD-B* associated with the Monte Amarelo flank collapse, and a transitional zone where the size of the hummocks decreases with distance from the source. The lateral extent of *MTD-B* may extend further south and west of the surveyed area. Dashed black line: previous mapped extent of the Monte Amarelo deposits (Masson *et al.* 2008); Grey line: eastern margin of the upper part of *MTD-B*. Solid black lines: location of the Parasound lines shown in (b) to (e). (b-e) Parasound profiles highlighting the acoustic character of (b) *MTD-A*, (c) the transitional zone, and (d-e) *MTD-B*. See the text for discussion on the nature of these deposits.

**Fig. 4.** (a) Compilation of sediment gravity cores retrieved south of Fogo and Santiago during M155 showing preliminary stratigraphic correlations based on visual analysis. Dashed red lines: correlations between different volcanoclastic sand layers (V1-V7); Dashed blue lines: correlation of background sediment layers (reddish-brown horizons) that helped to correlate the volcanoclastic sand layers. The location of these cores is shown in Figure 1c. (b) Zoom of gravity core M155\_38 (62-168 cm) showing the visually identified volcanoclastic layers V1-3.

**Fig. 5.** (a) Multibeam data overlain on GEBCO2019 bathymetry highlighting the presence of multiple fault-related escarpments and volcanic features on the shallow slopes south of Fogo. Additional mass-transport deposits are also identified in the Parasound data and coloured according to their burial depth (see legend). Solid red lines: location of the Parasound profiles

shown in (c) and (d). **(b)** Multibeam data showing the nature of the fault-related escarpments on the slopes south of Fogo and Santiago in more detail. **(c)** Parasound profile crossing one of the most striking escarpments visible on the western side (see 5b), highlights the relation between tectonic processes and landsliding in the area. Monte Amarelo *MTD-B* deposits (eastern side) are remobilised across the escarpment. **(d)** Parasound profile crossing an escarpment in the southern reaches of the working area highlights acoustically transparent MTDs at varying depths.

**Fig. 6.** *Southern Scour Complex (SSC)*. **(a)** Multibeam data highlighting the surface morphology of the *Southern Scour Complex* and the extension of the *Eastern Scour* (dashed white line). The location of Figure 6a with respect to the other features discussed in this article is shown in Figure 5. **(b-d)** Parasound profiles crossing the *SSC* showing a complex relationship between former and more recent landsliding processes, along with the variable acoustic character of the deposits and of the adjacent unfailed material. **(e)** Conceptual model showing the proposed retrogressive development of the *SSC* towards the north and east. Numbers 1-3 highlight the sequence of events.













